

RESEARCH PAPER

The checklist of Indian Phycitinae (Lepidoptera: Pyraloidea: Pyralidae)

SHIVASHARANARADDY¹, M. SHANKARAMURTHY², A. PRABURAJ, SOMASEKAR AND D. BASAVARAJA³

¹Department of Agricultural Entomology, College of Agriculture, Raichur

²Department of Agricultural Entomology, College of Agriculture, Bheemarayanagudi

³Department of Environmental sciences, College of Agriculture, Bheemarayanagudi
University of Agricultural Sciences, Raichur - 584 104, Karnataka, India

E-mail: smurthyent@gmail.com

(Received: June, 2020 ; Accepted: December, 2020)

Abstract: The subfamily Phycitinae was erected by Zeller in 1839 with type genus *Phycidea* Zeller, 1839. It is one of the economically important group of Pyraloidea, represents 3159 described species under 679 genera all over the world. In the present paper, a checklist was prepared for the species of Phycitinae occurring in India in a systematic sequence. As per the present checklist prepared, 233 species under 94 genera were documented. The report accounted for 7.38 per cent of species and 13.84 per cent of genera from India as compared to global record. The genera and species were listed alphabetically. Further, the valid name, type genus, type species, type locality and synonyms of each species under each genus were provided. The study provides base for further taxonomic research on this economically important group.

Key words: Phycitinae, Synonyms, Taxonomic, Type genus

Introduction

The Phycitinae was established by Zeller in 1839 with type genus *Phycidea*. It can be easily differentiated from other subfamilies of Pyraloidea by the absence of R_s vein in adults, the frenula of male and female adults' wings which contains a single bristle composed of several acanthi and in larvae, the mesothorax having sclerotised (hardened) ring around the base of seta. It is the second largest subfamily of Pyraloidea, comprises of 3159 described species under 679 genera all over the world (www.pyraloidea.org). Phycitinae is one of the economically important group of Pyraloidea, comprises of crop pests (borers and leaf webbers) and storage pests. Besides pests, it also includes predators and a weed killer.

Checklist is a skeletal classification of groups listed by the taxa for a quick reference. It provides a synopsis for the species occurring in certain region and also gives the current status of that particular species. In India, no efforts were made to prepare a checklist for Indian Phycitinae. However, Shankaramurthy *et al.* (2015) made a first effort to prepare a preliminary checklist of agriculturally important Pyraloidea of India, but they have included only 21 species of Phycitinae under 19 genera. Similarly, Nagaharish (2016) provided the checklist for Pyraloidea associated with horticultural crops in zone 1 and 2 of Karnataka, but he has included only 3 species of Phycitinae. In this context, an effort was made to provide the comprehensive checklist for the species of Phycitinae occurring in India which provides base for further taxonomic research on this economically important group.

Material and methods

The information for checklist of Phycitinae was collected from various primary and secondary sources of publications. The present study mainly referred the historic works of Hampson (1896); Munroe (1967); Arora (2000) and online web source (www.pyraloidea.org, 2020).

The present paper included all the species of Phycitinae occurring in India till date with comprehensive information on valid name, synonyms, type genus type species and type locality.

Results and discussion

The checklist prepared in the present paper for the Indian Phycitinae, included 233 species classified under 94 genera under the family Pyralidae. The synoptic checklist of Indian Phycitinae is provided here.

Subfamily: PHYCITINAE Zeller, 1839; type genus: *Phycidea* Zeller, 1839

Genus: *Acrobasis* Zeller, 1839 (*Nephopteryx*); type species: *Tinea consociella* Hübner, 1813

- = *Acrocaula* Hulst, 1900; type species: *Acrocaula comacornella* Hulst, 1900
- = *Catacrobasis* Gozmány, 1958; type species: *Tinea obtusella* Hübner, 1796,
- = *Conobathra* Meyrick, 1886; type species: *Conobathra automorpha* Meyrick, 1886
- = *Cyphita* Roesler, 1971; type species: *Myelois rufofusellus* Caradja, 1931
- = *Cyprusia* Amsel, 1958; type species: *Cyprusia wiltshirei* Amsel, 1958
- = *Mineola* Hulst, 1890; type species: *Myelois indigenella* Zeller, 1848
- = *Numonia* Ragonot, 1893; type species: *Numonia cymindella* Ragonot, 1893
- = *Seneca* Hulst, 1890; type species: *Cateremna tumidulella* Ragonot, 1887

- = *Trachycera* Ragonot, 1893; type species: *Rhodophaea pallicornella* Ragonot, 1887
- = *Hylopylora* Meyrick, 1933; type species: *Hylopylora craterantis* Meyrick, 1933
- = *Rhodophaeopsis* Amsel, 1950; type species: *Rhodophaea iranalis* Amsel, 1950
- Acrobasis atelogramma*** Meyrick, 1937 (***Myelois***); type locality: India (Uttar Pradesh: Gorakhpur)
- Genus: *Ammatucha*** Turner, 1922; type species: *Ammatucha lathria* Turner, 1922
- = *Umatraphycis* Roesler & Küppers, 1979; type species: *Nephopteryx semiirrorella* Hampson, 1896
- Ammatucha semiirrorella*** (Hampson, 1896) (***Nephopteryx***); type locality: India (Assam) & Bhutan
- Genus: *Ancova*** Ragonot, 1893; type species: *Nephopteryx meridionalis* Walker, 1863
- Ancova meridionalis*** Walker, 1863 (***Nephopteryx***); type locality: India
- Genus: *Ancylosis*** Zeller, 1839; type species: *Pyralis dilutella* Hübner sensu Treitschke, 1832
- = *Cabotia* Ragonot, 1888; type species: *Cabotia semidiscella* Ragonot, 1888
- = *Canarsiana* Strand, 1920; type species: *Canarsiana discocellularis* Strand, 1920
- = *Encystia* Hampson, 1901; type species: *Encystia bonhoti* Hampson, 1901
- = *Harnocha* Dyar, 1914; type species: *Harnocha velessa* Dyar, 1914
- = *Hulstia* Ragonot & Hampson, 1901; type species: *Nephopteryx undulatella* Clemens, 1860
- = *Gregorempista* Roesler, 1969; type species: *Nephopteryx validella* Christoph, 1877
- = *Heterographis* Ragonot, 1885; type species: *Euzophera samaritanella* Zeller, 1867
- = *Cornigerula* Amsel, 1935; type species: *Acornigerula bilineella* Amsel, 1935
- = *Mona* Hulst, 1888; type species: *Mona olbiella* Hulst, 1888
- = *Hypographia* Ragonot, 1890; type species: *Hypographia uncinatella* Ragonot, 1890
- = *Niethammeriodes* Roesler, 1969; type species: *Ancylosis diremptella* Ragonot, 1887
- = *Pseudocabotia* A. Blanchard & Knudson, 1985; type species: *Pseudocabotia balconiensis* A. Blanchard & Knudson, 1985
- = *Staudingeria* Ragonot, 1887; type species: *Ancylosis morbosella* Staudinger, 1879
- = *Cornigerula* Amsel, 1935; type species: *Cornigerula eremicola* Amsel, 1935
- = *Homoeograptia* Hampson, 1930; type species: *Staudingeria spectrifasciella* Ragonot, 1887
- = *Syria* Ragonot, 1887; type species: *Anerastia arenosella* Staudinger, 1859
- = *Hedemannia* Ragonot, 1887; type species: *Hedemannia l ineatella* Ragonot, 1887
- = *Iransharia* Amsel, 1959; type species: *Iransharia nigripunctella* Amsel, 1959
- Sub Genus: *Staudingeria*** Ragonot, 1887; type species: *Ancylosis morbosella* Staudinger, 1879
- = *Orniger.rapta* Hampson, 1930; type species: *Staudingeria spectrifasciella* Ragonot, 1887
- Ancylosis microstictella*** Hampson, 1903; type locality: Gujarat
- = *Heterographis microstictella* Hampson, 1903; type locality: India (Gujarat)
- Ancylosis rubripictella*** Hampson, 1901; type locality: Punjab
- = *Heterographis rubripictella* Hampson, 1901; type locality: India (Punjab)
- Genus: *Ancylodes*** Ragonot, 1887; type species: *Ancylodes pallens* Ragonot, 1887
- = *Heterographis rubripictella* Hampson, 1896; type locality: India (Punjab)
- Ancylodes argentescens*** (Hampson, 1919) (***Prosmixis***); type locality: South Africa
- Ancylodes arenacella*** Amsel, 1968; type locality: Pakistan (Karachi: Sindh)
- Ancylodes faustinella*** (Zeller, 1867) (***Euzophera***); type locality: Plains of Jordan
- = *Acornigerula bilineella* Amsel, 1935; type locality: Israel (South of Dead Sea)
- = *Heterographis airensis* Rothschild, 1921; type locality: Niger (Mountains of Baguezan: Asben)
- = *Heterographis damascenella* Amsel, 1952; type locality: Syria (Damascus)
- = *Heterographis rubripictella* Hampson, 1896; type locality: India (Punjab)
- = *Myelois staudingeri* Christoph, 1877; type locality: Iran (Schahrud)
- = *Staudingeria variabilis* Rothschild, 1915; type locality: Algeria
- Ancylodes illella*** Swinhoe, 1884; type locality: Pakistan (Karachi)
- Ancylodes karachiella*** (Amsel, 1968) (***Heterographis***); type locality: Pakistan (Karachi: Sindh)
- Ancylodes lapsalis*** (Walker, 1859) (Dosara); type locality: Sri Lanka

- Ancylodes mysorella* (Ragonot, 1888) (*Heterographis*); type locality: Indies
- Ancylodes microstictella* Hampson, 1903; type locality: India (Gujarat)
- Ancylodes niveicostella* Hampson, 1896; type locality: Sri Lanka
- Ancylodes obscuralis* (Hampson, 1912) (*Lygropia*); type locality: Venezuela (Cucuta)
- Ancylodes pakistanella* (Amsel, 1968) (*Heterographis*); type locality: Pakistan (Karachi: Sindh Province)
- Ancylodes trilineatella* Hampson, 1896; type locality: Pakistan (Punjab Province & Lahore)
- Ancylodes umbrilimbella* (Ragonot & Hampson, 1901) (*Heterographis*); type locality: Pakistan (Punjab Province & Multan)
- = *Heterographis concavella* Amsel, 1949; type locality: Iran (Ahvaz)
- = *Heterographis holella* D. Lucas, 1909: 71; type locality: Tunisia
- = *Heterographis sfaxella* D. Lucas, 1949: 96; type locality: Tunisia
- = *Heterographis thirtieta* D. Lucas, 1937; type locality: Algeria (Ouargla Province: Touggou)
- Genus:** *Anerastia* Hübner, 1825; type species: *Tinea lotella* Hübner, 1813
- = *Prinanerastia* Hampson, 1918; type species: *Tinea lotella* Hübner, 1813
- Anerastia celsella* (Walker, 1863) (*Araxis*); type locality: Sri Lanka
- Anerastia stictella* (Hampson, 1908) (*Anerastia*); type locality: Unknown
- Genus:** *Anonaepestis* Ragonot, 1894; type species: *Anonaepestis bengalella* Ragonot, 1894
- Anonaepestis bengalella* Ragonot, 1894; type locality: India & Bangladesh
- Genus:** *Anoristia* Ragonot, 1887; type species: *Anoristia umbrifasciella* Ragonot, 1887
- = *Ingridiola* Roesler, 1969; type species: *Heterographis conchyliaella* Ragonot, 1887
- Anoristia venosella* Moore, 1878 (*Homeosoma*); type locality: China
- Genus:** *Aphycita* Amsel, 1968; type species: *Aphycita sindella* Amsel, 1968
- Aphycita sindella* Amsel, 1968; type locality: Pakistan (Karachi)
- Genus:** *Apomyelois* Heinrich, 1956; type species: *Dioryctria bistriatella* Hulst, 1887
- Apomyelois ceratoniae* (Zeller, 1839) (*Myelois*); type locality: Austria (Laibach)
- = *Euzophera zellerella* Sorhagen, 1881; type locality: unknown
- = *Heterographis rivularis* Warren & Rothschild, 1905; type locality: Sudan (Atbara River: Nakheila)
- = *Hypsipyla psarella* Hampson, 1903; type locality: India
- = *Laodamia durandi* D. Lucas, 1950; type locality: Tunisia
- = *Myelois oporedestella* Dyar, 1911; type locality: USA (Florida: Miami)
- = *Myelois phoenicis* Durrant, 1915; type locality: Algeria
- = *Myelois tuerkheimiella* Sorhagen, 1881; type locality: unknown
- = *Phycis ceratoniella* Fischer von Röslerstamm, 1839
- = *Phycita dentilinella* Hampson, 1896; type locality: North East India (Manipur)
- = *Trachonitis pryrella* Vaughan, 1870; type locality: England (London)
- Apomyelois dentilinella* Hampson, 1896; type locality: North East India (Manipur)
- Apomyelois psarella* Hampson, 1903; type locality: India
- Genus:** *Aprophthasia* Amsel, 1968; type species: *Aprophthasia vartianae* Amsel, 1968
- Aprophthasia vartianae* Amsel, 1968; type locality: Pakistan (Karachi)
- Genus:** *Asarta* Zeller, 1848; type species: *Phycis aethiopella* Duponchel, 1837
- = *Chionea* Guenée, 1845; type species: *Phycis aethiopella* Duponchel, 1837
- Asarta alpicolella* Zeller, 1839; type locality: Switzerland
- = *Asarta alticola* Hampson, 1930 (*Anousterunia*); type locality: India (Kashmir: Rupshu)
- Genus:** *Asemeia* Hampson, 1930; type species: *Asemeia aprepi* Hampson, 1930
- Asemeia aprepi* Hampson, 1930; type locality: Unknown
- Genus:** *Assara* Walker, 1863; type species: *Assara albicostalis* Walker, 1863
- = *Cateremna* Meyrick, 1882; type species: *Euzophera microdoxa* Meyrick, 1882
- Assara albicostalis* Walker, 1863; type locality: Malaysia (Sarawak)
- Assara pinivora* Meyrick, 1933 (*Cateremna*); type locality: India (Kashmir: Hutmara)
- Assara tuberculosa* (Meyrick, 1933) (*Cateremna*); type locality: Myanmar (Burma)

Genus: *Auchmera* Hampson, 1930; type species:
Heterographis falsalis Hampson, 1908

Auchmera falsalis Hampson, 1908; type locality: Unknown

Genus: *Cadra* Walker, 1864; type species: *Cadra defectella* Walker, 1864

= *Xenephestia* Gozmány, 1958; type species: *Pempelia cautella* Walker, 1863

Cadra cautella (Walker, 1863) (*Pempelia*); type locality: Sri Lanka

Genus: *Caina* Ragonot, 1893; type species: *Caina deletella* Ragonot, 1893

Caina ciniferella Ragonot, 1893; type locality: India, Telangana State, Hyderabad

Caina deletella Ragonot, 1893; type locality: India (Maharashtra & Pune)

Caina inanitella (Ragonot, 1888) (*Anoristia*); type locality: India (Uttar Pradesh: Mainpuri)

Caina micrella Ragonot, 1893; type locality: Pakistan: Multan; India: Bombay; Pune

Genus: *Calguia* Walker, 1863; type species:
Calguia defiguralis Walker, 1863

= *Cabragus* Moore, 1886; type species: *Cabragus auritipalpus* Moore, 1886

= *Sigmarthria* Ragonot, 1888; type species: *Sigmarthria palpella* Ragonot, 1888

Calguia defiguralis Walker, 1863; type locality: Borneo

= *Cabragus auritipalpus* Moore, 1886

= *Nephopteryx bistriella* Pagenstecher, 1900

= *Salebria crocogastra* Meyrick, 1934; type locality: Indonesia (Java: Telawa)

= *Sigmarthria palpella* Ragonot, 1888; type locality: Malaysia (Sarawak)

Genus: *Canthelea* Walker, 1866; type species: *Homeosoma gratella* Walker, 1866

= *Gabra* Walker, 1866; type species: *Gabra tinealella* Walker, 1866.

Canthelea oegnusalis Walker, 1859 (*Pyrallis*); type locality: Srilanka

= *Epicrocis koshunalis* Strand, 1918; type locality: Taiwan

= *Eurhodope lateritialis* Walker, 1863; type locality: Srilanka

= *Homeosoma derasella* Swinhoe, 1885; type locality: India (Pune and Mumbai)

= *Homeosoma gratella* Walker, 1863; type locality: Srilanka

= *Myelois stibiella* Snellen, 1872; type locality: Guinea

= *Nephopteryx concorella* Walker, 1866; type locality: India

= *Nephopteryx laxalis* Swinhoe, 1885; type locality: India (Pune)

Genus: *Catastia* Hübner, 1825; type species: *Noctua marginea* Denis & Schiffermüller, 1775

= *Diosia* Duponchel, 1832; type species: *Noctua marginea* Denis & Schiffermüller, 1775

Catastia uniformalis Hampson, 1912; type locality: Indonesia (Sula Mangoli)

Genus: *Cathyalia* Ragonot, 1888; type species:
Cathyalia fulvella Ragonot, 1888

Cathyalia fulvella Ragonot, 1888; type locality: Indonesia (Sulawesi: Makassar)

= *Spatulipalpia erythrina* Hampson, 1896; type locality: India (Kolkata)

Genus: *Caustella* Hampson, 1930; type species:
Heterographis micralis Hampson, 1896

Caustella micralis Hampson, 1896 (*Heterographis*); type locality: India, Sri Lanka & Myanmar

Genus: *Cavipalpia* Ragonot, 1893; type species:
Cavipalpia translucidella Ragonot, 1893

Cavipalpia translucidella Ragonot, 1893; type locality: India (Hyderabad)

Genus: *Ceroprepes* Zeller, 1867; type species:
Ceroprepes patriciella Zeller, 1867

Ceroprepes patriciella Zeller, 1867; type locality: East India (Bihar, Jharkand, West Bengal and Odhisa)

Ceroprepes pulvillella (Zeller, 1867) (*Nephopteryx*); type locality: Eastern India (Bihar, Jharkand, West Bengal and Odhisa)

Ceroprepes ulvillella (Zeller, 1867) (*Nephopteryx*); type locality: Eastern India (Bihar, Jharkand, West Bengal and Odhisa)

Genus: *Citripestis* Ragonot, 1893; type species: *Nephopteryx sagittiferella* Moore, 1891

= *Philotroctis* Meyrick, 1933; type species: *Philotroctis eutrapphera* Meyrick, 1933

Citripestis sagittiferella (Moore, 1891) (*Nephopteryx*); type locality: Malaysia (Perak)

Citripestis pectinicornella (Hampson, 1896) (*Myelois*); type locality: Bhután

Citripestis eutrapphera (Meyrick 1933); type locality: Indonesia

Genus: *Comorta* Ragonot, 1888; type species:
Comorta atricostella Ragonot, 1888

Comorta atricostella Ragonot, 1888; type locality: Indies

Comorta nigracostalis Walker, 1863; type locality: Unknown

Genus: *Conobathra* Meyrick, 1886; type species:
Conobathra automorpha Meyrick, 1886

- = *Cyphita* Roesler, 1971; type species: *Myelois rufofusellus* Caradja, 1931
- = *Cyprusia* Amsel, 1958; type species: *Cyprusia wiltshirei* Amsel, 1958
- = *Mineola* Hulst, 1890; type species: *Myelois indigenella* Zeller, 1848
- = *Numonia* Ragonot, 1893; type species: *Numonia cymindella* Ragonot, 1893
- = *Styphlorachis* Hampson, 1930; type species: *Styphlorachis mesophaea* Hampson, 1930
- = *Seneca* Hulst, 1890; type species: *Cateremna tumidulella* Ragonot, 1887
- = *Trachycera* Ragonot, 1893; type species: *Rhodophaea pallicornella* Ragonot, 1887
- = *Hylopylora* Meyrick, 1933; type species: *Hylopylora craterantis* Meyrick, 1933
- = *Rhodophaeopsis* Amsel, 1950; type species: *Rhodophaea iranalis* Amsel, 1950

Conobathra aphidivora (Meyrick, 1934); type locality: Indonesia

- = *Cryptoblabes aphidivora* Yoshiyasu and Ohara, 1982; type locality: Japan

Conobathra atelogramma (Meyrick, 1937) (*Myelois*); type locality: India (Uttar Pradesh)

Genus: *Copamyntis* Meyrick, 1934; type species: *Elegia alectryonura* Meyrick, 1932

Copamyntis obliquifasciella (Hampson, 1896) (*Phycita*); type locality: India (HP)

Genus: *Crocidophora* Lederer, 1863; type species:
Crocidophora tubercularis Lederer, 1863

- = *Monocrocis* Warren, 1895; type species: *Hapalia flavofasciata* Moore, 1888

Crocidophora phaealis Hampson, 1903; type locality: India (Maharashtra & Mumbai)

Genus: *Cryptoblabes* Zeller, 1848; type species: *Ancylosis rutillella* Zeller, 1839

- = *Albinia* Briosi, 1877; type species: *Albinia wockiana* Briosi, 1877

Cryptoblabes proleucella Hampson, 1896; type locality: Sri Lanka

Cryptoblabes angustipennella Ragonot, 1888; type locality: Indies & India (Dharmasala)

Cryptoblabes ephestialis Hampson, 1903; type locality: Sri Lanka

Cryptoblabes flavizonalis Hampson, 1917; type locality: Indonesia

Cryptoblabes lophopterella Hampson, 1899; type locality: India (Meghalaya: Khasi Hills)

Cryptoblabes gnidiella Milliere, 1867; type locality: Spain, Barcelona.

Cryptoblabes proleucella Hampson, 1896; type locality: Sri Lanka

Genus: *Ctenomedes* Meyrick, 1935; type species:
Ctenomedes neuractis Meyrick, 1935

Ctenomedes neuractis Meyrick, 1935; type locality: India (Dehradun)

Genus: *Ctenomeristis* Meyrick, 1929; type species: *Ceroprepes almella* Meyrick, 1879

- = *Eremographa* Meyrick, 1932; type species: *Ceroprepes sebasmia* Meyrick, 1887

Ctenomeristis ebriola Meyrick, 1934; type locality: Sri Lanka (Gampola)

Genus: *Cyanaphycis* Roesler, 1978; type species:
Cyanaphycis bradleyella Roesler, 1978

Cyanaphycis bradleyella (Roesler, 1978) (*Cyanaphycis*); type locality: Pakistan (Kahuta)

Genus: *Didia* Ragonot, 1893; type species: *Didia subramosella* Ragonot, 1893

Didia strigivenella Hampson, 1891 (*Myelois*); type locality: India (Meghalaya, Khasi hills)

Genus: *Dioryctria* Zeller, 1846; typespecies: *Tinea abietella* (Denis and Schiffermüller), 1775

- = *Dioryctriodes* Mutuura & Munroe, 1974; type species: *Dioryctriodes daelei* Mutuura & Munroe, 1974

- = *Ocrisia* Ragonot, 1893; type species: *Myelois robiniella* Milliere, 1865

- = *Pinipestis* Grote, 1878; type species: *Nephopteryx zimmermanni* Grote, 1878

Dioryctria assamensis Mutuura, 1971; type locality: India (Assam, Upper Shillong)

Dioryctria castanea Bradley, 1969; type locality: India (Assam: Um Japung)

Dioryctria raoi Mutuura, 1971; type locality: North India, Almora, Ranikhet

Dioryctria symphoniella Hampson, 1899; type locality: India, Meghalaya, Khasi Hill

Genus: *Dipha* Arakaki & Yoshiyasu, 1988; type species:
Melitene aphidivora Meyrick, 1838

Dipha aphidivora (Meyrick, 1934) (*Melitene*); type locality: Indonesia (Java: Bogor)

Genus: *Eccopidia* Hampson, 1899; type species:
Eccopidia strigata Hampson, 1899

Eccopidia strigata Hampson, 1899; type locality: Unknown

Genus: *Elegia* Ragonot, 1887; type species:
Elegia atrifasciella Ragonot, 1887

= *Ichorarchis* Meyrick, 1937; type species: *Ichorarchis iozona* Meyrick, 1937

= *Microthrix* Ragonot, 1888; type species: *Microthrix fuscidorsella* Ragonot, 1888

Genus: *Emmalocera* Ragonot, 1888; type species:
Emmalocera crenatella Ragonot, 1888

= *Ambala* Ragonot, 1888; type species: *Ambala fuscostrigella* Ragonot, 1888

= *Baroda* Ragonot, 1888; type species: *Baroda paucigraphella* Ragonot, 1888

= *Critonia* Ragonot, 1891; type species: *Critonia subconcinella* Ragonot, 1891

= *Enosima* Ragonot in Ragonot & Hampson, 1901; type species: *Enosima neesimella* Ragonot, 1901

= *Lodiana* Ragonot, 1888; type species: *Lodiana umbrivittella* Ragonot, 1888

= *Papua* Ragonot, 1890; type species: *Papua latilimbella* Ragonot, 1890

= *Pectinigeria* Ragonot, 1888; type species: *Pectinigeria macrella* Ragonot, 1888

= *Poujadia* Ragonot, 1888; type species: *Poujadia sepicostella* Ragonot, 1888

= *Rhinaphe* Berg, 1875; type species: *Rhinaphe signicollis* Berg, 1875

= *Ampycophora* Meyrick, 1882; type species: *Pempelia apotomella* Meyrick, 1879

= *Singhaliella* Strand, 1920; type species: *Critonia simplicipalpis* Strand, 1920

= *Socora* Ragonot, 1888; type species: *Socora tenuicostella* Ragonot, 1888

Emmalocera apotomella (Meyrick, 1879) (*Pempelia*); type locality: Unknown

Emmalocera atricostella Ragonot, 1888; type locality: Indies

Emmalocera endopyrella Hampson, 1930; type locality: Mexico (Vera Cruz)

= *Synothmia bahamasella* Hampson, 1930; type locality: Bahamas

Emmalocera fuscostrigella (Ragonot, 1888) (*Ambala*); type locality: Indies

Emmalocera nigricostalis Walker, 1863; type locality: Unknown

Emmalocera ochracealis Hampson, 1912; type locality: Unknown

Emmalocera paucigraphella (Ragonot, 1888) (*Baroda*); type locality: Indies

Emmalocera umbricostella Ragonot, 1888; type locality: Indies

Emmalocera umbrivittella (Ragonot, 1888) (*Lodiana*); type locality: Indies

Genus: *Endolasia* Hampson, 1896; type species:
Endolasia melanoleuca Hampson, 1896

Endolasia melanoleuca Hampson, 1896; type locality: Unknown

= *Triaenoneura albifascia* Rebel, 1912; type locality: Egypt

Genus: *Ephestia* Guenée, 1845; type species: *Tinea elutella* Hübner, 1796

= *Anagasta* Heinrich, 1956; type species: *Ephestia kuehniella* Zeller, 1879

= *Hyphantidium* Scott, 1859; type species: *Hyphantidium sericarium* Scott, 1859

Ephestia glycyphloeas (Meyrick, 1935) (*Ephestia*); type locality: Iran

Ephestia rubrimediella Hampson, 1896; type locality: India (Meghalaya: Khashi Hills)

Genus: *Ephestiopsis* Ragonot, 1893; type species: *Myelois oenobarella* Meyrick, 1879

Ephestiopsis bipunctalis (Hampson, 1896) (*Cryptoblabe*); type locality: Bhutan

Genus: *Epicrocis* Zeller, 1848; type species:
Epicrocis festivella Zeller, 1848

= *Canthelea* Walker, 1866; type species: *Homoeosoma gratella* Walker, 1866

Epicrocis festivella Zeller, 1848; type locality: Indonesia (Java)

= *Gabra tinealella* Walker, 1866; type locality: Java

= *Scopula albigeralis* Walker, 1866; type locality: South Africa

= *Epicrocis varii* Balinsky, 1991; type locality: South Africa (Durban)

Epicrocis uberalis (Swinhoe, 1884) (*Pyrallis*); type locality: Pakistan (Karachi)

= *Candiope uberalis bahrainalis* Amsel, 1968; type locality: Bahrain (Rifaa Desert)

= *Susia uberalis desertella* (Caradja, 1910) (*Pristophora uberalis* var); type locality: Algeria (Biskra)

= *Bazaria fulvofasciata* Rothschild, 1915; type locality: Algeria (Oued Abbou)

= *Christophia cyrenaika* Turati, 1930; type locality: Libya (Cyrenaica: Augila)

= *Heterographis sahariensis* Rothschild, 1921; type locality: Nigeria (Damagarim: Makochia)

= *Pristophora discomaculella* Pauli f. obscura Turati, 1929; type locality: Libya (Sidi Mesri)

Genus: *Epischnia* Hübner, 1825; type species: *Tinea prodromella* Hübner, 1799

= *Epischnopsis* Amsel, 1954; type species: *Epischnia oculatella* Ragonot, 1887

Genus: *Ereboenis* Meyrick, 1935; type species: *Ereboenis saturata* Meyrick, 1935

Ereboenis saturata Meyrick, 1935; type locality: South India (Travancore)

Genus: *Ernophthora* Meyrick, 1887; type species: *Ernophthora phoenicias* Meyrick, 1887

= *Aspithra* Ragonot, 1888; type species: *Aspithra maculicostella* Ragonot, 1888

Ernophthora actiosoides Hampson, 1896; type locality: Unknown

Genus: *Etiella* Zeller, 1839; type species: *Phycis zinckenella* Treitschke, 1832

= *Alata* Walker, 1863; type species: *Alata anticalis* Walker, 1863

= *Arucha* Walker, 1863; type species: *Arucha indicatalis* Walker, 1863

= *Ceratamma* Butler, 1881; type species: *Ceratamma hobsoni* Butler, 1881

= *Mella* Walker, 1859; type species: *Mella* Walker, 1859

= *Modiana* Walker, 1863; type species: *Modiana dymnusalis scitivittalis* Walker, 1863

= *Rhamphodes* Guenée, 1845; type species: *Phycis etiella* Treitschke, 1835

Etiella zinckenella (Treitschke, 1832) (*Phycis*); type locality: Italy

Genus: *Euzophera* Zeller, 1867; type species: *Myelois cinerosella* Zeller, 1839

= *Ahwazia* Amsel, 1949; type species: *Ahwazia albocostalis* Amsel, 1949

= *Cymbalorissa* Gozmány, 1958; type species: *Stenoptycha fuliginosella*

= *Longignathia* Roesler, 1965; type species: *Longignathia cornutella* Roesler, 1965

= *Melia* Heinemann, 1865; type species: *Myelois cinerosella* Zeller, 1839

= *Pistogenes* Meyrick, 1937; type species: *Pistogenes mercatrix* Meyrick, 1937

= *Quadremipista* Roesler, 1973; type species: *Euzophera ultimella* Roesler, 1973

= *Stenoptycha* Heinemann, 1865; type species: *Myelois cinerosella* Zeller, 1839

Euzophera albicostalis Hampson, 1903; type locality: India (Kashmir: Sindh Valley)

Euzophera cocciphaga Hampson, 1908; type locality: India (Sikkim: Darjeeling)

Euzophera perticella Ragonot, 1888; type locality: India (Kolkata)

= *Euzophera nilghirisella* Ragonot, 1893; type locality: India (Kolkata)

Euzophera plumbeifasciella (Hampson, 1996) (*Euzophera*); type locality: India (West Bengal & Kolkatta)

Euzophera verrucicola (Hampson, 1896) (*Heterographis*); type locality: Sri Lanka

Genus: *Euzopherodes* Hampson, 1899; type species: *Euzopherodes albicans* Hampson, 1899

= *Delattinia* Roesler, 1964; type species: *Ephestia vapidella* Mann, 1857

= *Epilydia* Amsel, 1954; type species: *Myelois liturosella* Erschoff, 1874

= *Euzopherodes* Ragonot & Hampson 1901; type species: *Euzopherodes albicans*

= *Infinita* Whalley, 1970; type species: *Myelois lutisignella* Mann, 1869

= *Lydia* Rebel, 1901; type species: *Myelois lutisignella* Mann, 1869

= *Lydia* Ragonot & Hampson in Ragonot & Hampson, 1901; type species: *Myelois lutisignella* Mann, 1869

= *Neononia* Hampson, 1930; type species: *Nonia taprobalis* Hampson, 1908

= *Nyctigenes* Meyrick, 1937; type species: *Nyctigenes euphrontis* Meyrick, 1937

= *Phloeophaga* Chrétien, 1911; type species: *Euzophera pusilla* Mabilie, 1906

= *Radiestra* Hampson, 1927; type species: *Euzopherodes albistrigella* Hampson, 1908

= *Symphestia* Hampson, 1930; type species: *Euzopherodes ephestialis* Hampson, 1903

= *Rigonopyralis* Amsel, 1935; type species: *Trigonopyralis keltella* Amsel, 1935

Euzopherodes albicans Hampson, 1899; type locality: India (Kolkata) & Sri Lanka

= *albicans* Ragonot, 1901; type locality: India (Kolkatta) & Sri Lanka

Euzopherodes albistrigella Hampson, 1908; type locality: Unknown

Euzopherodes dendrophaga Meyrick, 1934; type locality: India (Kerala & Wyand)

Euzopherodes ephestialis Hampson, 1903; type locality: Unknown

Euzopherodes hemiphaealis (Hampson, 1907) (*Bradina*); type locality: Kenya

Euzopherodes proleucalis Hampson, 1908; type locality: India (Punjab)

Euzopherodes taprobalis Hampson, 1908; type locality: Unknown

Genus: *Gymnancyloides* Amsel, 1968; type species: *Gymnancyloides psorosella* Amsel, 1968

Gymnancyloides psorosella Amsel, 1968; type locality: Pakistan (Karachi)

Genus: *Homodigma* Hampson, 1930; type species: *Homodigma geera* Hampson, 1930

Homodigma geera Hampson, 1930; type locality: Unknown

Genus: *Homoeosoma* J. Curtis, 1833; type species: *Phycis gemina* Haworth, 1811

= *Anhomoeosoma* Roesler, 1965; type species: *Phycis nimbella* Duponchel, 1837

= *Ectohomoeosoma* Roesler, 1965; type species: *Ectohomoeosoma kasyellum* Roesler, 1965

= *Phycidea* Zeller, 1839; type species: *Tinea sinuella* Fabricius, 1794

= *Lotria* Guenée, 1845; type species: *Tinea sinuella* Fabricius, 1794

Homoeosoma ephestidiella Hampson, 1896; type locality: India (Nilgiris)

Homoeosoma goliathella Ragonot, 1888; type locality: Indies

Homoeosoma punctistrigella Ragonot, 1888; type locality: Pakistan (Punjab Province)

Genus: *Hydaspia* Ragonot, 1888; type species: *Hydaspia dorsipunctella* Ragonot, 1888

Hydaspia dorsipunctella Ragonot, 1888; type locality: Indies

Genus: *Hypargyria* Ragonot, 1888; type species: *Hypargyria metalliferella* Ragonot, 1888

Hypargyria metalliferella Ragonot, 1888; type locality: Indies & Ghana (Accra)

Genus: *Hypsipyla* Ragonot, 1888; type species: *Hypsipyla pagodella* Ragonot, 1888

Hypsipyla elachistalis Hampson, 1903; type locality: Sri Lanka

Hypsipyla robusta (Moore, 1886) (*Magiria*); type locality: Sri Lanka

= *Epicrocis terebrans* Olliff, 1890; type locality: Australia (New South Wales)

= *Hypsipyla pagodella* Ragonot, 1888; type locality: Indes

= *Hypsipyla scabrusculella* Ragonot, 1893; type locality: Madagascar

Hypsipyla rotundipex Hampson, 1903; type locality: India (Sikkim)

Genus: *Hypsotropa* Zeller, 1848; type species: *Hypsotropa limbella* Zeller, 1848

= *Tiarra* Ragonot, 1888; type species: *Tiarra pusillella* Ragonot, 1888

Genus: *Idiobrotis* Meyrick, 1937; type species: *Idiobrotis oxygrapha* Meyrick, 1937

Idiobrotis oxygrapha Meyrick, 1937; type locality: India (Tamil Nadu, Chennai & Coimbatore)

Genus: *Indomalayia* Roesler & Kuppers, 1979; type species: *Spatulipalpia flabellifera* Hampson, 1896

Indomalayia flabellifera (Hampson, 1896) (*Spatulipalpia*); type locality: Sri Lanka

= *Phycita umbratalis* Hampson, 1912; type locality: India

Indomalayia eugraphella Ragonot, 1888; type locality: Indies

= *Salebria iriditis* Meyrick, 1933; type locality: Indonesia (Java: Seneng)

= *Salebria pleurosaris* Meyrick, 1932; type locality: Indonesia (Java: Seneng)

Genus: *Kaurava* Roesler & Küppers, 1981; type species: *Rhodophaea rufimarginella* Hampson, 1896

Kaurava rufimarginella Hampson, 1896 (*Rhodophaea*); type locality: Bhutan

Genus: *Lacipea* Walker, 1863; type species: *Lacipea muscosella* Walker, 1863

Lacipea muscosella Walker, 1863; type locality: Unknown

Genus: *Letoa* Walker, 1866; type species: *Letoa patulella* Walker, 1866

Letoa patulella Walker, 1866; type locality: Unknown

Genus: *Mascelia* Hampson, 1930; type species: *Euzophera ectophoea* Hampson, 1908

Mascelia ectophoea Hampson, 1908; type locality: Unknown

Genus: *Merulempista* Roesler, 1967; type species: *Pempelia cingillella* Zeller, 1846

Merulempista cyclogramma Hampson, 1896 (*Phycita* (*Dioryctria*)); type locality: India (Assam)

Genus: *Mesciniodes* Hampson, 1901; type species: *Mesciniodes subinfractalis* Hampson, 1901

= *Mesciniella* Hampson, 1930; type species: *Euzophera micans* Hampson, 1896

Genus: *Metallosticha* Rebel, 1901; type species: *Myelois argyrogrammos* Zeller, 1847

= *Ametallosticha* Amsel, 1935; type species: *Ametallosticha ignore* Amsel, 1935

= *Metallosticha* Ragonot & Hampson, 1901; type species: *Myelois argyrogrammos* Zeller, 1847

- Metallosticha plumbeifasciella*** (Hampson, 1996) (***Euzophera***); type locality: India (West Bengal)
- Genus:** ***Metallostichodes*** Roesler, 1967; type species: ***Myelois nigrocyaneella*** Constant, 1865
- Metallostichodes hemicautea*** Hampson, 1899; type locality: Sri Lanka
- Genus:** ***Myelois*** Hübner, 1825; type species: ***Tinea cribrella*** Hübner, 1796
- = *Gnathogutta* Roesler, 1988; type species: *Myelois pumicosa* Lederer, 1855
- = *Lispe* Treitschke, 1832; type species: *Tinea cribrella* Hübner, 1796
- Myelois albicostalis*** Hampson, 1903; type locality: India (Kashmir: Shind Valley)
- Myelois echinopisella*** Chrétien, 1911; type locality: Algeria (Biskra)
- = *Myelois micropunctella* Amsel, 1951; type locality: Iran
- = *Myelois octopunctella* Amsel, 1968; type locality: Pakistan (Karachi)
- Myelois ephestialis*** Hampson, 1903; type locality: Unknown
- Myelois pectinicornella*** (Hampson, 1896) (***Myelois***); type locality: Bhutan
- Genus:** ***Nephopteryx*** Hübner, 1825; type species: ***Tinea angustella*** Hübner, 1796
- = *Alispa* Zeller, 1848; type species: *Tinea angustella* Hübner, 1796
- Nephopteryx acrobassella*** (Rebel, 1926); type locality: Egypt (Ramleh)
- = *Acrobassopsis talhouki* Amsel, 1958; type locality: Saudi Arabia
- Nephopteryx albifascialis*** (Hampson, 1917) (***Picrogama***); type locality: Papua New Guinea
- Nephopteryx asbolalis*** (Hampson, 1903) (***Nephopteryx***); type locality: India (Shimla)
- Nephopteryx eugraphella*** Ragonot, 1888; type locality: Indies
- Nephopteryx hemiargyralis*** (Hampson, 1908) (***Nephopteryx***); type locality: Unknown
- Nephopteryx lateritialis*** Walker, 1863; type locality: Sri Lanka
- Nephopteryx margarophanes*** Meyrick, 1932; type locality: Sri Lanka (Peradeniya)
- Nephopteryx rhyparella*** Ragonot, 1888; type locality: India (Punjab)
- = *Acrobasis nigrescens* Warren, 1888; type locality: India (Uttarakhand: Kalapani)
- Nephopteryx rubralis*** Hampson, 1900; type locality: Syria
- = *Aglossa pinguinalis f. lateritialis* Turati, 1921; type locality: Libya (Cyrénaique: Derna)
- = *Aglossa pinguinalis f. lividialis* Turati, 1921; type locality: Libya (Cyrénaique: Derna)
- Nephopteryx tumidella*** Zincken, 1818; type locality: Germany
- = *Phycita tumidana* Stephens, 1834; type locality: England
- Nephopteryx oedilepia*** Hampson, 1930; type locality: Unknown
- Nephopteryx striginervella*** Hampson, 1903; type locality: Unknown
- Genus:** ***Pempelia*** Hübner, 1825; type species: ***Tinea palumbella*** Denis & Schiffermüller, 1775
- = *Hoeneia* Caradja, 1938; type species: *Hoeneia sinensis* Caradja, 1938
- = *Salebria* Zeller, 1846; type species: *Tinea palumbella* Denis & Schiffermüller, 1775
- Pempelia heringii*** (Ragonot, 1889) (***Rhodophaea***); type locality: Sri Lanka
- = *Nephopteryx rubrizonella* Ragonot, 1893; type locality: Japan
- Pempelia palumbella*** (Denis & Schiffermüller, 1775) (***Tinea***); type locality: Austria
- = *Acrobasis undulatella* Walker, 1866; type locality: Sri Lanka
- = *Homoeosoma cinerea* Humphreys & Westwood, 1841; type locality: Great Britain
- Genus:** ***Phycita*** J. Curtis, 1828; type species: ***Tinea spissicella*** Fabricius, 1794
- = *Phycis* Fabricius, 1798; type species: *Tinea spissicella* Fabricius, 1794
- Phycita clientella*** Zeller, 1867: (***Nephopteryx***); type locality: India (Kolkatta)
- Phycita clientulella*** Ragonot, 1893; type locality: India (Bombay)
- Phycita eulepidella*** (Hampson, 1896: 94) (***Phycita***); type locality: Sri Lanka
- = *jasminophaga* Hampson, 1896; type locality: Unknown
- = *recondita* Turner, 1904; type locality: Australia
- = *similis* Whalley, 1962; type locality: Mozambique (Rennell Island)
- Phycita infusella*** (Meyrick, 1879 (***Nephopteryx***); type locality: Australia (Queensland)
- Phycita jasminophaga*** Hampson, 1896
- = *recondita* Turner, 1904; type locality: Australia

- = *similis* Whalley, 1962: 101; type locality: Mozambique (Rennell Island)
- Phycita nodicornella*** (Ragonot, 1888) (***Oligochroa***); type locality: Indies
- Phycita endochralis*** Hampson, 1908; type locality: Sri Lanka
- Phycita pachylepidella*** Ragonot & Hampson, 1901; type locality: Australia (Queensland)
- Phycita phaeella*** (Hampson, 1903: 25) (***Sandrabatis***); type locality: India (Sikkim)
- Phycita steniella*** Ragonot, 1890; type locality: Myanmar
- Phycita umbratalis*** Hampson, 1912; type locality: India
- Genus: *Phycitodes*** Hampson, 1917; type species: ***Phycitodes albistriata*** Hampson, 1917
- = *Rotruda* Heinrich, 1956; type species: *Homoeosoma mucidella* Ragonot, 1887
- Phycitodes albatella*** (Ragonot, 1887) (***Homoeosoma***); type locality: Uzbekistan (Samarkand)
- = *Phycitodes albatella dierli* Roesler, 1973; type locality: Nepal
- = *Homoeosoma parvum* Gerasimov, 1930; type locality: Uzbekistan (Bucharu)
- = *Phycitodes albatella pseudonimbella* (Bentinck, 1937) (***Homoeosoma***); type locality: Italy
- = *Phycitodes albatella reliquella* (Dyar, 1904) (***Homoeosoma***); type locality: USA
- Phycitodes albovittella*** (Ragonot, 1888) (***Homoeosoma***); type locality: India (Punjab)
- = *Homoeosoma costalbella* Amsel, 1954; type locality: Iraq (Basra)
- Phycitodes nepalensis*** Roesler, 1973; type locality: Nepal
- Genus: *Plodia*** Guenée, 1845; type species: ***Tinea interpunctella*** Hübner, 1796
- Plodia interpunctella*** (Hübner, 1810) (***Tinea***); type locality: Europe
- = *Unadilla latercula* Hampson, 1901; type locality: Bahamas (Nassau)
- Genus: *Polyocha*** Zeller, 1848; type species: ***Hypochalcia sanguinariella*** Zeller, 1848
- = *Epidauria* Rebel, 1901; type species: *Anerastia transversariella* Zeller, 1848
- = *Epidauria* Ragonot in Ragonot & Hampson, 1901; type species: *Anerastia transversariella* Zeller, 1848
- Polyocha depressellus*** (Swinhoe, 1885) (***Melissoblaptēs***); type locality: India (Pune)
- Genus: *Pristarthria*** Ragonot, 1893; type species: ***Salebria minutella*** Ragonot, 1885
- = *Sclerobia* Ragonot, 1893; type species: *Hypochalcia tritalis* Walker, 1863
- Pristarthria akbarella*** (Ragonot, 1888) (***Oligochroa***); type locality: Indies
- = *Spermatophthora pulverulenta* Warren, 1888; type locality: Pakistan (Punjab Province : Attock)
- Pristarthria mundalis*** (Walker, 1863) (***Nephopteryx***); type locality: Indonesia
- = *Oligochroa soricinella* Ragonot, 1888; type locality: Indies
- Genus: *Prorophora*** Ragonot, 1887; type species: ***Prorophora curvibasella*** Ragonot, 1887
- = *Proceratia* Amsel, 1950; type species: *Proceratia rhexogramma* Meyrick, 1937
- = *Reisserempista* Roesler, 1970; type species: *Prorophora mongolica* Roesler, 1970
- Prorophora dialeuca*** Hampson, 1912; type locality: Sri Lanka
- Prorophora afghanella*** Roesler, 1973; type locality: Pakistan
- Genus: *Protoetiella*** Inoue, 1959; type species: ***Protoetiella bipunctella*** Inoue, 1959
- = *Ketambeia* Roesler & Küppers, 1979; type species: *Etiella venustella* Hampson, 1896
- Protoetiella venustella*** (Hampson, 1896) (***Etiella***); type locality: Bhutan
- Genus: *Pseudodavara*** Roesler & Küppers, 1979; type species: ***Spatulipalpia haemaphoralis*** Hampson, 1908
- Pseudodavara haemaphoralis*** (Hampson, 1908) (***Spatulipalpia***); type locality: Sri Lanka
- Genus: *Psorosa*** Zeller, 1846; type species: ***Phycis dahliella*** Treitschke, 1832
- = *Ectyposa* Joannis, 1929; type species: *Phycis dahliella* Treitschke, 1832
- = *Sopsora* Roesler, 1975; type species: *Psorosa taishanella* Roesler, 1975
- Psorosa proleucalis*** Hampson, 1908; type locality: India (Punjab)
- Genus: *Ptyomaxia*** Hampson, 1903; type species: ***Ptyomaxia trigonifera*** Hampson, 1903
- = *Palibothra* Ragonot, 1890; type species: *Palibothra fuscogrisella* Ragonot, 1890
- Genus: *Raphimetopus*** Hampson, 1918; type species: ***Anerastia spinifrontella*** Ragonot, 1888
- Raphimetopus bimaculella*** (Ragonot, 1888) (***Anerastia***); type locality: Indies
- Raphimetopus spinifrontella*** (Ragonot, 1888) (***Anerastia***); type locality: Indies

Raphimetopus stigmatella (Ragonot, 1888) (*Anerastia*); type locality: Indies

Genus: *Rhodophaea* Guenée, 1845; type species: *Phycis dubiella* Duponchel, 1836

= *Aurana* Walker, 1863; type species: *Aurana actiosella* Walker, 1863

= *Longiculcita* Roesler, 1975; type species: *Rhodophaea vinaceella* Inoue, 1963

Rhodophaea albirenalis Hampson, 1908; type locality: Unknown

Rhodophaea nigralbella Hampson, 1899; type locality: India (Meghalaya: Khasis)

Genus: *Sandrabatis* Ragonot, 1893; type species: *Sandrabatis crassiella* Ragonot, 1893

Sandrabatis phaeella (Hampson, 1903) (*Sandrabatis*); type locality: India (Sikkim)

Sandrabatis crassiella Ragonot, 1893; type locality: India (Assam)

Genus: *Singhalia* Hampson, 1899; type species: *Citronia sarcoglaucia* Hampson, 1896

Singhalia cymogramma Meyrick, 1933; type locality: India (Mysore & Madras)

Singhalia sarcoglaucia Hampson, 1896; type locality: Unknown

Genus: *Spatulipalpia* Ragonot, 1893; type species: *Spatulipalpia effosella* Ragonot, 1893

Spatulipalpia albicostalis Hampson, 1912; type locality: Sri Lanka

Spatulipalpia albiatrialis Hampson, 1912; type locality: Sri Lanka

Spatulipalpia effosella Ragonot, 1893; type locality: India (Assam)

Spatulipalpia palliostalis (Walker, 1963) (*Nephopteryx*); type locality: India

Genus: *Taprobania* Hampson, 1930; type species: *Homoeosoma glaucochroa* Hampson, 1908

Taprobania glaucochroa Hampson, 1908; type locality: Unknown

Genus: *Tephris* Ragonot, 1890; type species: *Pempelia cyriella* Erschoff, 1874

= *Aphyletes* Ragonot, 1893; type species: *Salebria nigrisparsella* Ragonot, 1887

Tephris connexella Ragonot, 1893; type locality: India (Hyderabad)

Tephris perlucidella Ragonot, 1893; type locality: Pakistan

Genus: *Thermopteryx* Hampson, 1912; type species: *Thermopteryx rubrifusa* Hampson, 1912

Thermopteryx rubrifusa Hampson, 1912; type locality: Unknown

Genus: *Thiallela* Walker, 1863; type species: *Thiallela signifera* Walker, 1863

= *Luconia* Ragonot, 1888; type species: *Luconia pallidobasella* Ragonot, 1888

= *Phalobathra* Meyrick, 1932; type species: *Phalobathra escigera* Meyrick, 1932

Thiallela endochralis Hampson, 1908; type locality: Sri Lanka

Thiallela ligeralis (Walker, 1863) (*Myelois*); type locality: Sri Lanka

= *Luconia pallidobasella* Ragonot, 1888; type locality: Malaysia (Sarawak)

= *Thiallela signifera* Walker, 1863; type locality: Malaysia (Sarawak)

Genus: *Thylacoptila* Meyrick, 1885; type species: *Thylacoptila paurosema* Meyrick, 1885

= *Bussa* Ragonot, 1888; type species: *Bussa maculella* Ragonot, 1888

Thylacoptila paurosema Meyrick, 1885; type locality: Portugal (Cape Verde, São Vicente)

= *Bussa maculella* Ragonot, 1888; type locality: Ghana (Accra)

= *Nephopteryx canescentella* Hampson, 1896; type locality: Burma (Thayetmyo)

Genus: *Tinestra* Hampson, 1908; type species: *Tinestra micralis* Hampson, 1908

Tinestra micralis Hampson, 1908; type locality: Unknown

Genus: *Trissonca* Meyrick, 1882; type species: *Spermatophthora mesactella* Meyrick, 1879

Trissonca oxyopa Meyrick, 1934; type locality: India (Rajasthan: Thar Desert)

Genus: *Vinicia* Ragonot, 1893; type species: *Salebria gypsopa* Meyrick, 1883

= *Eleusina* Ragonot, 1901; type species: *Ephestia guttella* Snellen, 1882

Vinicia phloeophaga (Meyrick, 1935) (*Eleusina*); type locality: Pakistan

Genus: *Volobilis* Walker, 1863; type species: *Volobilis biplaga* Walker, 1863

Volobilis chloropterella (Hampson, 1896) (*Phycita*); type locality: Sri Lanka & Myanmar Yangon & Borneo

= *Crambus decapitatus* Strand, 1918; type locality: Taiwan

Genus: *Zynodes* Whalley, 1970; type species: *Blabioides strigerella* Hampson, 1903

= *Blabioides* Hampson, 1903; type species: *Blabioides strigerella* Hampson, 1903

Zynodes strigerella (Hampson, 1903) (*Blabioides*); type locality: Sri Lanka

In the earlier studies, Shankaramurthy *et al.* (2015) prepared a checklist for 21 species of Phycitinae occurring in India. In another study, Nagaharish (2016) provided the checklist for 3 species of Phycitinae associated with horticultural crops in

zone 1 and 2 of Karnataka. In the current study, checklist was prepared for 233 species of Phycitinae occurring in India. The current study provides base for further taxonomic research on this economically important group.

Conclusion

The present study is the first consolidated (up to 2020) comprehensive checklist of all the species of Phycitinae occurring in India, which comprises about 233 described species under 94 genera. The study provides base for further taxonomic research

References

- Arora G S, 2000, Studies on some Indian pyralid species of economic importance, part-1, Crambinae, Schoenobinae, Nymphulinae, Phycitinae and Gallerinae (Lepidoptera: Pyraloidea). *Records of the Zoological Survey of India*, Calcutta. 181 (8): 1-169.
- Hampson G F, 1896, The fauna of British India, including Ceylon and Burma. Moths-Taylor and Francis Ltd., 4: 594.
- Munroe E G, 1967, Revision of North American species of *Udea* Guenée (Lepidoptera: Pyralidae). *Memoirs of the Entomological Society of Canada*, 49: 1-57.
- Nagaharish G, 2016, Faunistic studies on Pyralidae (Lepidoptera: Pyraloidea) associated with horticultural crops from zone-1 and 2 of Karnataka. *M. Sc. Thesis*, University of Agricultural Sciences, Raichur, Karnataka, India.
- Shankaramurthy M, Nagaraj S K and Prabhuraj A, 2015, Agriculturally important Pyraloidea (Lepidoptera) of India: key to subfamilies, current taxonomic status and a preliminary checklist. *Entomology*, 40 (1): 23-62.
- Zeller P C, 1839, Die knotenhornigen Phyciden nach ihren Arten beschrieben - Isis von Oken, *Leipzig*, 32 (3):167-219.
- www.pyraloidea.org, 2020